

No Dense Subgraphs Appear in the Triangle-free Graph Process

Stefanie Gerke Tamás Makai
 Royal Holloway College
 University of London
 Egham TW20 0EX

30.01.2010

Abstract

Consider the triangle-free graph process, which starts from the empty graph on n vertices and a random ordering of the possible $\binom{n}{2}$ edges; the edges are added in this ordering provided the graph remains triangle free. We will show that there exists a constant c such that no copy of any fixed finite triangle-free graph on k vertices with at least ck edges asymptotically almost surely appears in the triangle-free graph process.

Keywords: triangle-free graph process, subgraphs

1 Introduction

The random graph process starts from the empty graph on n vertices $G_{n,0}$ and in the i th step, $G_{n,i}$ is obtained from $G_{n,i-1}$ by inserting an edge chosen uniformly at random from $\overline{G_{n,i-1}}$, the complement graph of $G_{n,i-1}$. We are interested in structural properties of $G_{n,m}$ that have probability tending to one as the number of vertices n tends to infinity. We say that these properties hold asymptotically almost surely (a.a.s.). The random graph process is well understood, partly as $G_{n,m}$ has strong connections to the random graph model $G_{n,p}$ when $p \approx m/\binom{n}{2}$. In $G_{n,p}$ each edge is present independently of the presence or absence of all other edges with probability p ; see [2], [5].

We are interested in the triangle-free graph process where at step i of the random graph process an edge is chosen uniformly at random from the set

of edges which are not only in $\overline{G_{n,i-1}}$ but when added to $G_{n,i-1}$ the graph remains triangle free. The process terminates when no more edges can be inserted. In this paper we show that there exists a constant c such that a.a.s. no copy of any fixed finite triangle-free graph on k vertices with at least ck edges appears in the triangle-free graph process. For instance large complete bipartite graphs a.a.s. do not appear in the triangle-free graph process.

Wolfovitz[8] proved recently a complementary result namely that balanced sparse graphs appear in the triangle-free process. More precisely, he gave bounds on the number of copies of any fixed balanced triangle-free graph F with $e_F < 2v_F$ that a.a.s hold in the triangle-free graph process, after a small percentage of the edges have been inserted. Here, v_F denotes the number of vertices in F , e_F denotes the number of edges and a graph F is balanced if $e_H/v_H \leq e_F/v_F$ for every induced subgraph $H \subset F$. Let us note that a related process, the random planar graph process (where at each step an edge is inserted if the graph remains planar) behaves differently. Gerke, Schlatter, Steger and Taraz[4] showed that a.a.s. the planar graph process contains a copy of any fixed planar graph after inserting just $(1 + \varepsilon)n$ edges.

Erdős, Suen and Winkler[3] were the first to consider the triangle-free graph process. They have shown that the triangle-free graph process terminates a.a.s. after $O(n^{3/2}\sqrt{\log n})$ edges have been inserted in contrast to the more restrictive property of being bipartite, which a.a.s. terminates after $O(n^2)$ steps. More recently Bohman[1] strengthened this result by showing a conjecture of Spencer[7], namely that the final graph contains a.a.s. $\Theta(n^{3/2}\sqrt{\log n})$ edges. He also proved that the maximal independent set has a.a.s. size $O(\sqrt{n \log n})$. This implies Kim's result[6] on the lower bound of the Ramsey number $R(3, t) = \Omega(t^2 / \log t)$.

In his proof Bohman analyzed the steps leading to an edge being excluded from the triangle-free graph process until a small percentage of the edges have been inserted. For any pair of non-adjacent vertices u, v , he gave estimates on the number of “open”, “partial” and “complete” vertices see Figure 1. In particular, if there is a vertex which is complete with respect to $\{u, v\}$, then the edge $\{u, v\}$ cannot be inserted by the triangle-free process. In this case $\{u, v\}$ is called a closed pair.




Figure 1. w is open/partial/complete with respect to $\{u, v\}$

Using Bohman's estimates, we will show that in the regime when the estimates hold, a.a.s. no copy of a fixed dense subgraph F appears and more importantly, for any possible placement of a copy of F one of its edges becomes closed. Therefore no copy of F can be completed later in the process.

2 Main Results

Denote the graph created after i edges were inserted by the triangle-free random graph process with G_i and the set of its edges with E_i . A pair of vertices $\{u, v\}$ is called closed at step i if inserting the edge $\{u, v\}$ in G_i would result in a triangle. The set of all closed pairs at step i is denoted by C_i . A pair is open if it is neither an edge of the graph nor closed. The set of open pairs at step i is denoted by O_i and $Q(i) = |O_i|$. Note that both vertices and pairs of vertices can be open. If $\{u, v\} \notin E_i$ then $Y_{u,v}(i)$ is the set of partial vertices, that is, the set of vertices w such that exactly one of the pairs $\{u, w\}$ or $\{v, w\}$ is open and the other is an edge at step i :

$$Y_{u,v}(i) = \{w \in V : |\{\{u, w\}, \{v, w\}\} \cap O_i| = |\{\{u, w\}, \{v, w\}\} \cap E_i| = 1\}.$$

If $\{u, v\} \in E_i$ then $Y_{u,v}(i) = Y_{u,v}(i-1)$.

Define $t(i) = i/n^{3/2}$. Bohman[1] showed bounds on $|O_i|$ and $|Y_{u,v}(i)|$ for $i \leq \mu n^{3/2} \sqrt{\log n}$ with $\mu = 1/32$. (Bohman made no effort to optimize the value of μ .) For the remainder of the paper we set $\mu = 1/32$ and $m = \mu n^{3/2} \sqrt{\log n}$.

Definition 1. Let H be the event that the following bounds hold for all pairs $\{u, v\} \notin E_i$ and for all $i \leq m = \mu n^{3/2} \sqrt{\log n}$ with $\mu = 1/32$:

$$\begin{aligned} |Q(i) - n^2 q(t(i))| &\leq n^2 g_q(t(i)) \\ \left| |Y_{u,v}(i)| - \sqrt{n} y(t(i)) \right| &\leq \sqrt{n} g_y(t(i)) \end{aligned}$$

where

$$\begin{aligned}
q(t) &= \exp(-4t^2)/2 \\
y(t) &= 4t \exp(-4t^2) \\
g_q(t) &= \begin{cases} \exp(41t^2 + 40t)n^{-1/6} & : t \leq 1 \\ \frac{\exp(41t^2 + 40t)}{t}n^{-1/6} & : t > 1 \end{cases} \\
g_y(t) &= \exp(41t^2 + 40t)n^{-1/6}.
\end{aligned}$$

Theorem 1. [1] *The event H holds a.a.s..*

Note that $t(m) = \mu\sqrt{\log n}$. Let $W \subset V$ and define $e_i(W)$ as the number of edges spanned by W after step i .

Lemma 2. *Fix k . Let S_k be the event that there exists $W \subset V$ with $|W| = k$ and $e_m(W) \geq 3k$. Then $P(S_k|H) = o(1)$.*

Proof. Fix k vertices in G and denote this set by W . Let A_i be the event that an edge is added between two vertices in W at step i . Then

$$\begin{aligned}
P(A_i|H) &\leq \frac{k^2}{Q(i)} \leq \frac{k^2}{n^2(q(t(i)) - g_q(t(i)))} \leq \frac{2k^2}{n^2q(t(i))} \leq \frac{2k^2}{n^2q(t(m))} \\
&\leq \frac{4k^2}{n^2 \exp(-4\mu^2 \log n)} = \frac{4k^2}{n^{2-4\mu^2}}.
\end{aligned}$$

Therefore

$$\begin{aligned}
P(e_m(W) \geq 3k|H) &\leq \binom{m}{3k} \left(\frac{4k^2}{n^{2-4\mu^2}} \right)^{3k} \leq \left(\frac{em4k^2}{3kn^{2-4\mu^2}} \right)^{3k} \\
&\leq \left(\frac{e\mu n^{3/2} \sqrt{\log n} 4k}{3n^{2-4\mu^2}} \right)^{3k} = o\left(\frac{1}{n^{3k/2-20k\mu^2}} \right).
\end{aligned}$$

Since there are $\binom{n}{k}$ ways to select k vertices, it follows from the union bound that

$$P(S_k|H) \leq \binom{n}{k} o\left(\frac{1}{n^{3k/2-20k\mu^2}} \right) = o\left(\frac{n^k}{n^{3k/2-20k\mu^2}} \right) = o(1)$$

as μ^2 is sufficiently small. $\square$

Given a fixed graph F , we say that there exists a copy of F in G if a function $f : V(F) \rightarrow V(G)$ exists such that $\{f(u), f(v)\} \in E(G)$ for all $\{u, v\} \in E(F)$. We have just shown that no copy of a dense graph appears in the process while the first m edges are taken. We will now show that when m edges have been taken at least one edge of any copy of F is closed.

Theorem 3. *Let T be the event that there exists a copy of a graph F with e edges and k vertices satisfying $10k/\mu^2 \leq e$ in the triangle-free graph process. Then $P(T | [\overline{S_k}, \overline{S_{2k}}, H]) = o(1)$.*

Proof. Fix a set of vertices W with $|W| = k$, and a set of pairs of vertices $E_F \subset W \times W$ such that if the pairs in E_F were inserted as edges they would form a copy of F on W . Let $C_F(i)$ be the event that at least one pair in E_F is closed after step i and $O_F(i)$ be the event that none is closed after step i . For the following assume we are in the event $O_F(i)$.

Note that a pair $\{u, v\}$ is closed at step i if and only if there is a partial vertex $w \in Y_{u,v}(i)$ and the missing edge is chosen. Thus the probability of closing a pair $s \in O_i$ is $|Y_s(i)|/Q(i)$. The problem is that an edge can close several pairs of vertices. The subset of $W \times W$ closed by $\{w_j, v\} \in O_i$, with $v \notin W$ is $w_j \times (N_i(v) \cap W)$ (see Figure 2), where $N_i(v)$ denotes the neighbourhood of v in G_i .




Figure 2. The edge $\{v, w_1\}$ closes $\{w_1, w_2\}, \{w_1, w_3\}, \{w_1, w_4\}$

Let D_i be the set of vertices not in W that have more than 6 neighbours in W at time i . Excluding the pairs with both vertices in W and the pairs with a vertex in D_i the remaining pairs close at most 6 pairs in $W \times W$ and in particular at most 6 pairs in E_F . Therefore $\sum_{f \in E_F(W) \setminus E_i} |Y_f(i) \setminus (D_i \cup W)|$ counts any pair that closes a pair in E_F at most 6 times.

Since we are in $\overline{S_{2k}}$, the set D_i can have size at most k otherwise $W \cup D_i$ would span more than $6k$ edges. Hence

$$\begin{aligned}
P(C_F(i+1)|[O_F(i), \overline{S_k}, \overline{S_{2k}}, H]) &\geq \frac{\sum_{f \in E_F(W) \setminus E_i} |Y_f(i) \setminus (D_i \cup W)|}{6Q(i)} \\
&\geq \frac{\sum_{f \in (E_F) \setminus E_i} (|Y_f(i)| - 2k)}{6Q(i)}.
\end{aligned}$$

Since we are in the event $\overline{S_k}$ there are at most $3k$ edges in E_F also $|E_F| \geq 10k/\mu^2$ so the sum is over at least $(10/\mu^2 - 3)k \geq 9k/\mu^2$ open pairs. Thus

$$P(C_F(i+1)|[O_F(i), \overline{S_k}, \overline{S_{2k}}, H]) \geq \frac{9k}{\mu^2} \frac{\sqrt{n}(y(t(i)) - g_y(t(i))) - 2k}{6n^2(q(t(i)) + g_q(t(i)))}.$$

If n is large enough then $q(t(i)) + g_q(t(i)) \leq 2q(t(i))$, and for $m \geq i \geq n^{4/3}$ we have

$$y(t(i)) - g_y(t(i)) \geq \frac{y(t(i))}{2} \geq 2t(n^{4/3}) \exp(-4t^2(m)) = 2n^{-\frac{1}{6}-4\mu^2},$$

therefore since k is a constant:

$$\frac{\sqrt{n}y(t(i))}{2} - 2k \geq \frac{7}{15} \sqrt{n}y(t(i)),$$

and so :

$$\begin{aligned}
P(C_F(i+1)|[O_F(i), \overline{S_k}, \overline{S_{2k}}, H]) &\geq \frac{9k}{\mu^2} \frac{7\sqrt{n}y(t(i))/15}{6n^2(q(t(i)) + g_q(t(i)))} \\
&\geq \frac{7k}{\mu^2} \frac{\sqrt{n}y(t(i))}{20n^2q(t(i))} = \frac{7k}{\mu^2} \frac{4t(i) \exp(-4t^2(i))}{10n^{3/2} \exp(-4t^2(i))} \\
&= \frac{14ki}{5\mu^2 n^3}.
\end{aligned}$$

It follows that for $m \geq i \geq n^{4/3}$ and sufficiently large n ,

$$P(O_F(i+1)|[O_F(i), \overline{S_k}, \overline{S_{2k}}, H]) \leq 1 - \frac{14ki}{5\mu^2 n^3} \leq \exp\left(-\frac{14ki}{5\mu^2 n^3}\right).$$

Thus for sufficiently large n

$$\begin{aligned}
P(O_F(m)|[\overline{S_k}, \overline{S_{2k}}, H]) &= \prod_{i=0}^{m-1} P(O_F(i+1)|[O_F(i), \overline{S_k}, \overline{S_{2k}}, H]) \\
&\leq \prod_{i=\lceil n^{4/3} \rceil}^{m-1} \exp\left(-\frac{14ki}{5\mu^2 n^3}\right) = \exp\left(\sum_{i=\lceil n^{4/3} \rceil}^{m-1} -\frac{14ki}{5\mu^2 n^3}\right) \\
&= \exp\left(-\frac{14k}{5\mu^2 n^3} \left(\frac{m(m-1)}{2} - \frac{\lceil n^{4/3} \rceil(\lceil n^{4/3} \rceil - 1)}{2}\right)\right) \\
&\leq \exp\left(-\frac{4k}{3} \frac{m^2}{\mu^2 n^3}\right) = \exp\left(-\frac{4k}{3} \log n\right) = n^{-4k/3}.
\end{aligned}$$

Applying the union bound gives

$$P(T|[\overline{S_k}, \overline{S_{2k}}, H]) \leq \binom{n}{k} k! n^{-4k/3} \leq n^{k-4k/3} = o(1).$$

□

References

- [1] Tom Bohman. The triangle-free process. *Adv. Math.*, 221(5):1653–1677, 2009.
- [2] Béla Bollobás. *Random graphs*, volume 73 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2001.
- [3] Paul Erdős, Stephen Suen, and Peter Winkler. On the size of a random maximal graph. In *Proceedings of the Sixth International Seminar on Random Graphs and Probabilistic Methods in Combinatorics and Computer Science, “Random Graphs ’93” (Poznań, 1993)*, volume 6, pages 309–318, 1995.
- [4] Stefanie Gerke, Dirk Schlatter, Angelika Steger, and Anusch Taraz. The random planar graph process. *Random Structures Algorithms*, 32(2):236–261, 2008.
- [5] Svante Janson, Tomasz Łuczak, and Andrzej Ruciński. *Random graphs*. Wiley-Interscience Series in Discrete Mathematics and Optimization. Wiley-Interscience, New York, 2000.

- [6] Jeong Han Kim. The Ramsey number $R(3, t)$ has order of magnitude $t^2/\log t$. *Random Structures Algorithms*, 7(3):173–207, 1995.
- [7] Joel Spencer. Maximal triangle-free graphs and Ramsey $R(3, t)$. *Unpublished manuscript*, 1995.
- [8] Guy Wolfowitz. Triangle-free subgraphs at the triangle-free process. *preprint*.